

LQTS

LQTS

What is Long QT Syndrome (LQTS)?

LQTS is a condition that causes problems with the electrical activity of your heart. This is usually due to inheriting a faulty gene from your mother or your father. This results in the chemicals, sodium and potassium, that make the electricity in your heart, becoming out of balance. It takes longer for your heart to reset itself electrically after every beat and can show up on an electrocardiogram (ECG) as a measurement which is bigger than normal, the long QT interval. Your heart can then be more at risk of having abnormal heart rhythms that start in the bottom of your heart (the ventricles).

LQTS LONG QT SYNDROME

What are the symptoms?

Some people get more symptoms than others who may have none at all. The main, and most important symptom, is collapsing and blacking out suddenly. Occasionally you may feel palpitations where your heart seems to have extra beats or missing beats. There are usually no physical signs of LQTS. In some people LQTS can cause a cardiac arrest. This is when your heart stops pumping blood around the body and you stop breathing normally. It is very dangerous and resuscitation is needed.

Things to discuss in clinic

- List of any medicines you should avoid that can worsen the QT interval
- Loud noises/startle may cause blackouts in some people
- Physical activity can still continue but may need to be changed
- Pregnancy and genetic counselling
- Contraception
- Illegal drugs and alcohol can often make long QT worse
- Counselling/support groups

What tests will I need?

Tests include ECGs, 24-48 hour ECG monitoring and exercise ECG tests, which are all painless and involve having sticky labels on your chest. Often genetic testing is needed which can be done on a small amount of your blood or sometimes a small amount of your spit.

Some genes with faults have been found in Long QT syndrome. So even if you have Long QT syndrome it may be that your family gene isn't yet known. Or you may have a faulty gene that can cause Long QT syndrome but a long QT interval doesn't yet show on your ECG, so you may need to keep going to clinic for ECG tests in case it does show up.

All of this can be difficult to cope with, especially if you have had a sudden death in your family from Long QT syndrome.

What treatment will I need?

It depends on how high your risk is of getting an abnormal ventricular heart rhythm (from the bottom of your heart). Those at most risk are people who have had a cardiac arrest or blackouts/collapses, a very long QT on the ECG, girls in puberty and young adulthood and boys before puberty.

If you are at high risk, especially if you've had a cardiac arrest then an ICD is needed. You may also get beta blocker tablets which can be very protective. Sometimes a pacemaker is a good alternative to an ICD in younger children who are at higher risk. They may also have an operation on the left side of the neck to treat the nerves that go to the heart, called a sympathectomy.


If you are not so high risk then only beta blockers may be needed. Sometimes other drugs are used.

Please see our leaflets on ICDs and pacemakers if you are going to have one.


Thanks to Dr Behr for contributions to the text.

Normal electrical pathway

Find out more about your heart and visit:
c-r-y.org.uk
yheart.net/meet



Ventricular arrhythmia



QT interval



normal QT interval

prolonged/long QT interval

British Heart Foundation
 Greater London House
 180 Hampstead Road
 London NW1 7AW
 Phone: 020 7554 0000
 Fax: 020 7554 0100
 Website: bhf.org.uk
 Registered Charity Number in
 England and Wales (225971)
 and in Scotland (SC039426)

Cardiac Risk in the Young
 Unit 7, Epsom Downs
 Metro Centre, Waterfield,
 Tadworth, Surrey KT20 5LR
 Phone: 01737 363 222
 Fax: 01737 363 444
 Website: c-r-y.org.uk
 email: cry@c-r-y.org.uk
 Registered Charity Number in
 England and Wales 1050845